

Ocean City Public Schools Pre-Participation Forms



Student Name	Gender	Grade	Sport
HISTORY REVIEWED AND S	TUDENT EXAMINEI	BY:	Physician's/Provider's Stamp:
☐ Primary Care Provider			
☐ School Physician Provider			
☐ License Type:			
□ MD/DO			
\Box APN			
□ PA			
PHYSICIAN'S/PROVIDER'S SIGNAT	URE:		
Today's Date:	Date of Exam:		
	DIAN of the child listed above Registration and agree to al	ceancity s Regist ve, I con l guideli	ration" and follow instructions mpleted the online portion of ines and policies mentioned.
☐ <u>Use and Misuse of Opioid Drug</u>	s Fact Sheet		
☐ Opioid Education Video Proced	<u>ure</u>		
☐ Sports-Related Concussion and	Head Injury Fact Sheet		
☐ Sudden Cardiac Death in Young	g Athletes Pamphlet		
Parent/Guardian Name:			
Parent/Guardian Signature:			
Date:			
Please return this form along with the	other sign-off sheets listed	above t	o the above to the Health Office

by the first day of try-outs. Any student that does not return this form will not be allowed to participate -

Rev.01/18

- NO EXCEPTIONS!

■ PREPARTICIPATION PHYSICAL EVALUATION

HISTORY FORM

(Note: This form is to be filled out by the patient and parent prior to seeing the physician. The physician should keep a copy of this form in the chart.)

Date	e of Exam					
	ne					
Sex	Age Grade Sch	ool		Sport(s)		
Me	edicines and Allergies: Please list all of the prescription and over	-the-co	unter m	redicines and supplements (herbal and nutritional) that you are currently	taking	
	you have any allergies?	ntify spe	ecific all	lergy below. □ Food □ Stinging Insects		
Expl	lain "Yes" answers below. Circle questions you don't know the an	swers t	0.			
ė	NERAL QUESTIONS	Yes	No	MEDICAL QUESTIONS	Yes	No
1.	Has a doctor ever denied or restricted your participation in sports for any reason?			26. Do you cough, wheeze, or have difficulty breathing during or after exercise?		
2.	Do you have any ongoing medical conditions? If so, please identify below: ☐ Asthma ☐ Anemia ☐ Diabetes ☐ Infections Other:			27. Have you ever used an inhaler or taken asthma medicine? 28. Is there anyone in your family who has asthma?		
3.	Have you ever spent the night in the hospital?			29. Were you born without or are you missing a kidney, an eye, a testicle (males), your spleen, or any other organ?		
4.	Have you ever had surgery?			30. Do you have groin pain or a painful bulge or hernia in the groin area?		
HE	ART HEALTH QUESTIONS ABOUT YOU	Yes	No	31. Have you had infectious mononucleosis (mono) within the last month?		
5.	Have you ever passed out or nearly passed out DURING or AFTER exercise?			32. Do you have any rashes, pressure sores, or other skin problems? 33. Have you had a herpes or MRSA skin infection?		
6.	Have you ever had discomfort, pain, tightness, or pressure in your chest during exercise?			34. Have you ever had a head injury or concussion?		
7.	Does your heart ever race or skip beats (irregular beats) during exercise?			35. Have you ever had a hit or blow to the head that caused confusion, prolonged headache, or memory problems?		
8.	Has a doctor ever told you that you have any heart problems? If so,			36. Do you have a history of seizure disorder?		
	check all that apply: ☐ High blood pressure ☐ A heart murmur			37. Do you have headaches with exercise?		
	☐ High cholesterol ☐ A heart infection ☐ Kawasaki disease Other:			38. Have you ever had numbness, tingling, or weakness in your arms or legs after being hit or falling?		
9.	Has a doctor ever ordered a test for your heart? (For example, ECG/EKG, echocardiogram)			39. Have you ever been unable to move your arms or legs after being hit or falling?		
10.	Do you get lightheaded or feel more short of breath than expected			40. Have you ever become ill while exercising in the heat?		
L.	during exercise?			41. Do you get frequent muscle cramps when exercising?		
\vdash	Have you ever had an unexplained seizure?			42. Do you or someone in your family have sickle cell trait or disease?	\vdash	
12.	Do you get more tired or short of breath more quickly than your friends during exercise?			43. Have you had any problems with your eyes or vision?	\vdash	
HE	ART HEALTH QUESTIONS ABOUT YOUR FAMILY	Yes	No	44. Have you had any eye injuries? 45. Do you wear glasses or contact lenses?		
13.	Has any family member or relative died of heart problems or had an unexpected or unexplained sudden death before age 50 (including			46. Do you wear protective eyewear, such as goggles or a face shield? 47. Do you worry about your weight?		
14.	drowning, unexplained car accident, or sudden infant death syndrome)? Does anyone in your family have hypertrophic cardiomyopathy, Marfan syndrome, arrhythmogenic right ventricular cardiomyopathy, long QT			48. Are you trying to or has anyone recommended that you gain or lose weight?		
	syndrome, short QT syndrome, Brugada syndrome, or catecholaminergic			49. Are you on a special diet or do you avoid certain types of foods?		
	polymorphic ventricular tachycardia?			50. Have you ever had an eating disorder?		
15.	Does anyone in your family have a heart problem, pacemaker, or implanted defibrillator?			51. Do you have any concerns that you would like to discuss with a doctor?		
16.	Has anyone in your family had unexplained fainting, unexplained			FEMALES ONLY		
	seizures, or near drowning?			52. Have you ever had a menstrual period?	\vdash	
-	NE AND JOINT QUESTIONS	Yes	No	53. How old were you when you had your first menstrual period? 54. How many periods have you had in the last 12 months?	_	
17.	Have you ever had an injury to a bone, muscle, ligament, or tendon that caused you to miss a practice or a game?			Explain "yes" answers here		
18.	Have you ever had any broken or fractured bones or dislocated joints?			Lapiani yes answers nere		
19.	Have you ever had an injury that required x-rays, MRI, CT scan, injections, therapy, a brace, a cast, or crutches?					
20.	Have you ever had a stress fracture?] ————		
21.	Have you ever been told that you have or have you had an x-ray for neck instability or atlantoaxial instability? (Down syndrome or dwarfism)					
22.	Do you regularly use a brace, orthotics, or other assistive device?					
\vdash	Do you have a bone, muscle, or joint injury that bothers you?					
\vdash	Do any of your joints become painful, swollen, feel warm, or look red?					
25.	Do you have any history of juvenile arthritis or connective tissue disease?] —————		
	ereby state that, to the best of my knowledge, my answers to		•	·		
Signa	ature of athlete Signature o	ı parent/g	uardian _	Date		

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

HEUSOS

New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71

■ PREPARTICIPATION PHYSICAL EVALUATION

THE ATHLETE WITH SPECIAL NEEDS: SUPPLEMENTAL HISTORY FORM

Name	·			Date of birth		
Sex	Age	Grade	School	Sport(s)		
-	Type of disability Date of disability					
_	Classification (if available)					
_		and andidant/trauma ather)				
-	Cause of disability (birth, disea					
J. I	List the sports you are interest	leu III piayiliy			Yes	No
6 1	Do you regularly use a brace,	assistive device or prostheti	c?		100	110
_	Do you use any special brace					
-	Do you have any rashes, press					
-	Do you have a hearing loss? D					
_	10. Do you have a visual impairment?					
-	Do you use any special device		ion?			
12. I	Do you have burning or discor	nfort when urinating?				
13. I	Have you had autonomic dysre	eflexia?				
14. I	Have you ever been diagnosed	d with a heat-related (hypert	hermia) or cold-related (hypothermia) illne	ss?		
15. I	Do you have muscle spasticity	?				
16. I	Do you have frequent seizures	that cannot be controlled by	y medication?			
Expla	in "yes" answers here					
Pleas	e indicate if you have ever h	any of the following				
1000	o maioato n you mato otor n	iaa aiiy or alo lollolliligi			Yes	
						l No
Atlan	toaxial instability				103	No
_	toaxial instability y evaluation for atlantoaxial in	stability			103	NO
X-ray		stability			100	NO
X-ray Dislo	v evaluation for atlantoaxial in	stability			163	No
X-ray Dislo	y evaluation for atlantoaxial in cated joints (more than one)	stability			103	NO
X-ray Dislo	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen	stability			103	NO
X-ray Dislo Easy Enlar Hepa	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen	stability			103	No
X-ray Dislo Easy Enlar Hepa Oste	y evaluation for atlantoaxial in cated joints (more than one) bleeding rged spleen tititis	stability			100	No
X-ray Dislo Easy Enlar Hepa Oste	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen utitis openia or osteoporosis	stability			100	No
X-ray Dislo Easy Enlar Hepa Oste Diffic Num	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen utitis openia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or his	ands			100	No
X-ray Dislo Easy Enlar Hepa Oste Diffic Num Num	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen utitis openia or osteoporosis cutty controlling bowel cutty controlling bladder bness or tingling in arms or his bness or tingling in legs or fee	ands			100	No
X-ray Dislo Easy Enlar Hepa Oste Diffic Num Num Weal	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen utitis openia or osteoporosis cutty controlling bowel cutty controlling bladder bness or tingling in arms or habness or tingling in legs or feat kness in arms or hands	ands				No
X-ray Dislo Easy Enlar Hepa Oste Diffic Num Num Weal	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen utitis openia or osteoporosis cutty controlling bowel cutty controlling bladder bness or tingling in arms or habness or tingling in legs or fee kness in arms or hands kness in legs or feet	ands				No
X-ray Dislo Easyy Enlar Hepa Oste Diffic Num Num Weal Weal Rece	y evaluation for atlantoaxial in cated joints (more than one) bleeding gged spleen utitis openia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or his bness or tingling in legs or fee kness in arms or hands kness in legs or feet int change in coordination	ands				No
X-ray Dislot Easy Enlar Hepa Oste Diffic Num Num Weal Weal Rece	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen tittis openia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or his bness or tingling in legs or feek senses in arms or hands kness in legs or feet and tohange in coordination and change in ability to walk	ands				No
X-ray Dislot Easy Enlar Hepa Oste Diffic Num Num Weal Rece Rece Spin:	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen tittis openia or osteoporosis culty controlling bladder breas or tingling in arms or his bress or tingling in legs or feekness in arms or hands kness in legs or feet and change in coordination and change in coordination and change in ability to walk a bifida	ands				No
X-ray Dislot Easy Enlar Hepa Oste Diffic Num Num Weal Rece Rece Spin:	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen tittis openia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or his bness or tingling in legs or feek senses in arms or hands kness in legs or feet and tohange in coordination and change in ability to walk	ands				No
X-ray Dislot Easy Enlar Hepz Oste Diffic Num Num Weal Recce Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen tittis openia or osteoporosis culty controlling bladder breas or tingling in arms or his bress or tingling in legs or feekness in arms or hands kness in legs or feet and change in coordination and change in coordination and change in ability to walk a bifida	ands				No
X-ray Dislot Easy Enlar Hepz Oste Diffic Num Num Weal Recce Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding gred spleen tittis popenia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or hands kness in arms or hands kness in legs or feet and than the change in coordination at change in ability to walk a bifida k allergy	ands				NO
X-ray Dislo Easy Enlan Hepz Oste Diffic Num Num Weal Weal Rece Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding gred spleen tittis popenia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or hands kness in arms or hands kness in legs or feet and than the change in coordination at change in ability to walk a bifida k allergy	ands				No
X-ray Dislo Easy Enlan Hepz Oste Diffic Num Num Weal Weal Rece Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding gred spleen tittis popenia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or hands kness in arms or hands kness in legs or feet and than the change in coordination at change in ability to walk a bifida k allergy	ands				No
X-ray Dislo Easy Enlan Hepz Oste Diffic Num Num Weal Weal Rece Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding gred spleen tittis popenia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or hands kness in arms or hands kness in legs or feet and than the change in coordination at change in ability to walk a bifida k allergy	ands				No
X-ray Dislo Easy Enlan Hepz Oste Diffic Num Num Weal Weal Rece Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding gred spleen tittis popenia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or hands kness in arms or hands kness in legs or feet and than the change in coordination at change in ability to walk a bifida k allergy	ands				No
X-ray Dislo Easy Enlan Hepz Oste Diffic Num Num Weal Weal Rece Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding gred spleen tittis popenia or osteoporosis culty controlling bowel culty controlling bladder bness or tingling in arms or hands kness in arms or hands kness in legs or feet and than the change in coordination at change in ability to walk a bifida k allergy	ands				No
X-ray Dislote Easy Enlau Hepa Oste Diffic Num Num Weal Recce Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen tititis openia or osteoporosis culty controlling bowel culty controlling blowel bness or tingling in arms or his bness or tingling in legs or feek kness in legs or feet and that change in coordination and to thank a bifida a tallergy in "yes" answers here	ands et	rs to the above questions are complete	and correct.		No
X-ray Dislot Easy Enlan Hepa Oste Diffic Num Weal Weal Rece Spin: Late:	y evaluation for atlantoaxial in cated joints (more than one) bleeding ged spleen tititis openia or osteoporosis culty controlling bowel culty controlling blowel bness or tingling in arms or his bness or tingling in legs or feek kness in legs or feet and that change in coordination and to thank a bifida a tallergy in "yes" answers here	ands et	rs to the above questions are complete Signature of parent/quardian	and correct.	Date	No

■ PREPARTICIPATION PHYSICAL EVALUATION

PHYSICAL EXAMINATION FORM

Name			[Date of birth
PHYSICIAN REMINDERS				
1. Consider additional questions on more sensitive issues Do you feel stressed out or under a lot of pressure? Do you ever feel sad, hopeless, depressed, or anxious? Do you feel safe at your home or residence? Have you ever tried cigarettes, chewing tobacco, snuff, or dip? During the past 30 days, did you use chewing tobacco, snuff, or dip? Do you drink alcohol or use any other drugs? Have you ever taken anabolic steroids or used any other performanc have you ever taken any supplements to help you gain or lose weigh Do you wear a seat belt, use a helmet, and use condoms? Consider reviewing questions on cardiovascular symptoms (questions)	e supplement? t or improve your p	erformance?		
EXAMINATION	J-14).			
Height Weight	□ Male	☐ Female		
BP / (/) Pulse	Vision R	1 20/	L 20/	Corrected □ Y □ N
MEDICAL		NORMAL		ABNORMAL FINDINGS
Appearance • Marfan stigmata (kyphoscoliosis, high-arched palate, pectus excavatum, ar arm span > height, hyperlaxity, myopia, MVP, aortic insufficiency)	achnodactyly,			
Eyes/ears/nose/throat Pupils equal Hearing				
Lymph nodes				
Heart a • Murrunrs (auscultation standing, supine, +/- Valsalva) • Location of point of maximal impulse (PMI) Pulses				
Simultaneous femoral and radial pulses				
Lungs				
Abdomen				
Genitourinary (males only) ^b				
Skin HSV, lesions suggestive of MRSA, tinea corporis				
Neurologic °				
MUSCULOSKELETAL				
Neck				
Back				
Shoulder/arm				
Elbow/forearm Wrist/hand/fingers				
Hip/thigh				
Knee				
Leg/ankle				
Foot/toes				
Functional • Duck-walk, single leg hop				
*Consider ECG, echocardiogram, and referral to cardiology for abnormal cardiac history or ex *Consider GU exam if in private setting, Having third party present is recommended. *Consider cognitive evaluation or baseline neuropsychiatric testing if a history of significant of Cleared for all sports without restriction				
☐ Cleared for all sports without restriction with recommendations for further e	valuation or treatme	nt for		
□ Not cleared				
□ Pending further evaluation				
☐ For any sports				
☐ For certain sports				
Reason				
Recommendations				
I have examined the above-named student and completed the preparticip participate in the sport(s) as outlined above. A copy of the physical exam arise after the athlete has been cleared for participation, a physician may to the athlete (and parents/guardians).	is on record in my or rescind the clearan	office and can be ma	de available to t	he school at the request of the parents. If conditio
Name of physician, advanced practice pures (APN), physician assistant (DA) (marine) (h. m. e.)			Data

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

Signature of physician, APN, PA ____

■ PREPARTICIPATION PHYSICAL EVALUATION CLEARANCE FORM

Name	Sex 🗆 M 🗆 F Age	Date of birth
☐ Cleared for all sports without restriction		
☐ Cleared for all sports without restriction with recommendations for further e	evaluation or treatment for	
☐ Not cleared		
☐ Pending further evaluation		
☐ For any sports		
☐ For certain sports		
Reason		
Recommendations		
EMERGENCY INFORMATION		
Allergies		
Other information		
I have examined the above-named student and completed the pro- clinical contraindications to practice and participate in the sport(and can be made available to the school at the request of the par the physician may rescind the clearance until the problem is reso (and parents/guardians).	s) as outlined above. A copy of the phys ents. If conditions arise after the athlete	ical exam is on record in my office has been cleared for participation,
Name of physician, advanced practice nurse (APN), physician assistant (P	' A)	Date
Address		
Signature of physician, APN, PA		
Completed Cardiac Assessment Professional Development Module		
DateSignature		

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71



OPIOID USE AND MISUSE EDUCATIONAL FACT SHEET

Keeping Student-Athletes Safe

School athletics can serve an integral role in students' development. In addition to providing healthy forms of exercise, school athletics foster friendships and camaraderie, promote sportsmanship and fair play, and instill the value of competition.

Unfortunately, sports activities may also lead to injury and, in rare cases, result in pain that is severe or long-lasting enough to require a prescription opioid painkiller. It is important to understand that overdoses from opioids are on the rise and are killing Americans of all ages and backgrounds. Families and communities across the country are coping with the health, emotional and economic effects of this epidemic.²

This educational fact sheet, created by the New Jersey Department of Education as required by state law (*N.J.S.A.* 18A:40-41.10), provides information concerning the use and misuse of opioid drugs in the event that a health care provider prescribes a student-athlete or cheerleader an opioid for a sports-related injury. Student-athletes and cheerleaders participating in an interscholastic sports program (and their parent or guardian, if the student is under age 18) must provide their school district written acknowledgment of their receipt of this fact sheet.

How Do Athletes Obtain Opioids?

In some cases, student-athletes are prescribed these medications. According to research, about a third of young people studied obtained pills from their own previous prescriptions (i.e., an unfinished prescription used outside of a physician's supervision), and 83 percent of adolescents had unsupervised access to their prescription medications.³ It is important for parents to understand the possible hazard of having unsecured prescription medications in their households. Parents should also understand the importance of proper storage and disposal of medications, even if they believe their child would not engage in non-medical use or diversion of prescription medications.

What Are Signs of Opioid Use?

According to the National Council on Alcoholism and Drug Dependence, 12 percent of male athletes and 8 percent of female athletes had used prescription opioids in the 12-month period studied.³ In the early stages of abuse, the athlete may exhibit unprovoked nausea and/or vomiting. However, as he or she develops a tolerance to the drug, those signs will diminish. Constipation is not uncommon, but may not be reported. One of the most significant indications of a possible opioid addiction is an athlete's decrease in academic or athletic performance, or a lack of interest in his or her sport. If these warning signs are noticed, best practices call for the student to be referred to the appropriate professional for screening, 4 such as provided through an evidence-based practice to identify problematic use, abuse and dependence on illicit drugs (e.g., Screening, Brief Intervention, and Referral to Treatment (SBIRT)) offered through the New Jersey Department of Health.

According to NJSIAA Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies

indicate that about 80 percent of neroin users started out by abusing narcotic painkillers."

What Are Some Ways Opioid Use and Misuse Can Be Prevented?

According to the New Jersey State Interscholastic Athletic Association (NJSIAA) Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

The Sports Medical Advisory Committee, which includes representatives of NJSIAA member schools as well as experts in the field of healthcare and medicine, recommends the following:

- The pain from most sports-related injuries can be managed with non-narcotic medications such as acetaminophen, nonsteroidal anti-inflammatory medications like ibu profen, na proxen or aspirin. Read the label carefully and always take the recommended dose, or follow your doctor's instructions. More is not necessarily better when taking an over-the-counter (OTC) pain medication, and it can lead to dangerous side effects.
- Ice therapy can be utilized appropriately as an anesthetic.
- Always discuss with your physician exactly what is being prescribed for pain and request to avoid narcotics.
- In extreme cases, such as severe trauma or post-surgical pain, opioid pain medication should not be prescribed for more than five days at a time;
- Parents or guardians should always control the dispensing of pain medications and keep them in a safe, non-accessible location; and
- Unused medications should be disposed of immediately upon cessation of use. Ask your pharmacist about drop-off locations
 or home disposal kits like Deterra or Medsaway.






STATE OF NEW JERSEY DEPARTMENT OF HEALTH

NJSIAA SPORTS MEDICAL **ADVISORY COMMITTEE**





Even With Proper Training and Prevention, Sports Injuries May Occur

There are two kinds of sports injuries. Acute injuries happen suddenly, such as a sprained ankle or strained back. Chronic injuries may happen after someone plays a sport or exercises over a long period of time, even when applying overuse-preventative techniques.5

Athletes should be encouraged to speak up about injuries, coaches should be supported in injury-prevention decisions, and parents and young athletes are encouraged to become better educated about sports safety.6

What Are Some Ways to Reduce the Risk of Injury?

Half of all sports medicine injuries in children and teens are from overuse. An overuse injury is damage to a bone, muscle, ligament, or tendon caused by repetitive stress without allowing time for the body to heal. Children and teens are at increased risk for overuse injuries because growing bones are less resilient to stress. Also, young athletes may not know that certain symptoms are signs of overuse.

The best way to deal with sports injuries is to keep them from happening in the first place. Here are some recommendations to consider:



PREPARE Obtain the preparticipation physical evaluation prior to participation on a school-sponsored interscholastic or intramural athletic team or squad.



CONDITIONING Maintain a good fitness level during the season and offseason. Also important are proper warm-up and cooldown



PLAY SMART Try a variety of sports and consider specializing in one sport before late adolescence to help avoid overuse injuries.



ADEQUATE HYDRATION Keep the body hydrated to help the heart more easily pump blood to muscles, which helps muscles work efficiently.



TRAINING Increase weekly training time, mileage or repetitions no more than 10 percent per week. For example, if running 10 miles one week, increase to 11 miles the following week. Athletes should also cross-train and perform sport-specific drills in different ways, such as running in a swimming pool instead of only running on the road.



REST UP Take at least one day off per week from organized activity to recover physically and mentally. Athletes should take a combined three months off per year from a specific sport (may be divided throughout the year in one-month increments). Athletes may remain physically active during rest periods through alternative low-stress activities such as stretching, yoga or walking.



PROPER EQUIPMENT Wear appropriate and properly fitted protective equipment such as pads (neck, shoulder, elbow, chest, knee, and shin), helmets, mouthpieces, face quards, protective cups, and eyewear. Do not assume that protective gear will prevent all injuries while performing more dangerous or risky activities.

Resources for Parents and Students on Preventing Substance Misuse and Abuse

The following list provides some examples of resources:

National Council on Alcoholism and Drug Dependence - NJ promotes addiction treatment and recovery.

New Jersey Department of Human Services, Division of Mental Health and Addiction Services has a mission to decrease the abuse of alcohol, tobacco and other drugs by supporting the development of a comprehensive network of prevention, intervention and treatment services in New Jersey.

New Jersey Prevention Network includes a parent's quiz on the effects of opioids.

Operation Prevention Parent Toolkit is designed to help parents learn more about the opioid epidemic, recognize warning signs, and open lines of communication with their children and those in the community.

Parent to Parent NJ is a grassroots coalition for families and children struggling with alcohol and drug addiction.

Partnership for a Drug Free New Jersey is New Jersey's anti-drug alliance created to localize and strengthen drug-prevention media efforts to prevent unlawful drug use, especially among young people.

ReachNJ provides information for parents and families, including addiction and treatment stories.

The Science of Addiction: The Stories of Teens shares common misconceptions about opioids through the voices of teens.

Youth IMPACTing NJ is made up of youth representatives from coalitions across the state of New Jersey who have been impacting their communities and peers by spreading the word about the dangers of underage drinking, marijuana use, and other substance misuse.

- References 1 Massachusetts Technical Assistance Partnership for Prevention
 - ² Centers for Disease Control and Prevention New Jersey State Interscholastic Athletic
- Association (NJSIAA) Sports Medical Advisory Committee (SMAC)
- ⁴ Athletic Management, David Csillan, athletic trainer, Ewing High School, NJSIAA SMAC
- 5 National Institute of Arthritis and Musculoskeletal and Skin Diseases
- 6 USA TODAY
- 7 American Academy of Pediatrics

An online version of this fact sheet is available on the New Jersey Department of Education's Alcohol, Tobacco, and Other Drug Use webpage. Updated Jan. 23, 2018.

OCEAN CITY SCHOOL DISTRICT

501 Atlantic Avenue, Suite 1 Ocean City, New Jersey 08226 – 3891

Phone: (609) 399-5150 www.oceancityschools.org



Use and Misuse of Opioid Drugs Fact Sheet Student-Athlete and Parent/Guardian Sign-Off

In accordance with N.J.S.A. 18A:40-41.10, public school districts, approved private schools for students with disabilities, and nonpublic schools participating in an interscholastic sports program must distribute the <u>Opioid Use and Misuse Educational Fact Sheet</u> to all student-athletes and cheerleaders. In addition, schools and districts must obtain a signed acknowledgement of receipt of the fact sheet from each student-athlete and cheerleader, and for students under age 18, the parent or guardian must also sign.

This sign-off sheet is due to the Health Office prior to the first official practice session of every student-athlete or cheerleader. This acknowledgement is required before <u>each</u> season in which the student-athlete or cheerleader will be participating.

Name of School: Ocean City Intermediate School
Name of School District (if applicable): Ocean City School District
I/We acknowledge that we received and reviewed the online version of the Educational Fact Sheet on the Use and Misuse of Opioid Drugs while completing the online sign-up form for sports within the Ocean City School District.
Student Name:
Student Signature:
Parent/Guardian Signature:
Date:

NEW JERSEY STATE INTERSCHOLASTIC ATHLETIC ASSOCIATION

1161 Route 130 North, Robbinsville, NJ 08691 Phone 609-259-2776 ~ Fax 609-259-3047

Memorandum

To: All Athletic Directors of Member Schools

From: Tony Maselli, Assistant Director

Date: June 2019

Re: Opioid Education Video Procedure

To All Athletic Directors:

Acting to address the increased risk of opioid abuse among high school athletes, the Office of the New Jersey Coordinator for Addiction Responses and Enforcement Strategies (NJCARES) and the New Jersey State Interscholastic Athletic Association (NJSIAA) announced on February 19, 2019, a new partnership to educate student athletes and their parents/guardians on addiction risks associated with sports injuries and opioid use.

This educational initiative, spearheaded by Attorney General Gurbir Grewal and approved by the Executive Committee of the NJSIAA, is a collaborative effort to use video programming to raise awareness among high school athletes that they face a higher risk of becoming addicted to prescription pain medication than their fellow students who do not play sports.

Beginning with the 2019 fall season, we are making available to all student athletes and their parents/guardians, an educational video about the risks of opioid use as it relates to student athletes. The video will be available on August 1, 2019 and can be found on the NJSIAA website under "Athlete Wellness" which is located under the "Health & Safety tab. We are strongly encouraging student athletes and parents/guardians to watch the video as soon as it becomes available. An acknowledgement that students and their parents/guardians have watched the video will be required starting with the 2019-2020 winter season.

All member schools are asked to add to their current athletic consent forms the sign-off listed below. The sign-off acknowledgment is an NJSIAA mandate; student athletes are required to view the video only once per school year prior to the first official practice of the season in their respective sport, but the signed acknowledgment is required for each sport a student participates in. Athletes that are 18 years or older do not need the parents/guardians to watch the video.

Opioid Video is located at: https://youtu.be/3Rz6rkwpAx8

NJSIAA OPIOID POLICY ACKNOWLEDGEMENT

We have viewed the NJ CARES educational video on the risks of opioid use for high school athletes. We understand the NJSIAA policy that requires students, and their parents(s)/guardian(s) if a student is under the age of 18, to view this video and sign this acknowledgement.

Student's Signature:	Date:
Parent/Guardian Signature:	Date:

Sports-Related Concussion and Head Injury Fact Sheet and Parent/Guardian Acknowledgement Form

A concussion is a brain injury that can be caused by a blow to the head or body that disrupts normal functioning of the brain. Concussions are a type of Traumatic Brain Injury (TBI), which can range from mild to severe and can disrupt the way the brain normally functions. Concussions can cause significant and sustained neuropsychological impairment affecting problem solving, planning, memory, attention, concentration, and behavior.

The Centers for Disease Control and Prevention estimates that 300,000 concussions are sustained during sports related activities nationwide, and more than 62,000 concussions are sustained each year in high school contact sports. Second-impact syndrome occurs when a person sustains a second concussion while still experiencing symptoms of a previous concussion. It can lead to severe impairment and even death of the victim.

Legislation (P.L. 2010, Chapter 94) signed on December 7, 2010, mandated measures to be taken in order to ensure the safety of K-12 student-athletes involved in interscholastic sports in New Jersey. It is imperative that athletes, coaches, and parent/guardians are educated about the nature and treatment of sports related concussions and other head injuries. The legislation states that:

- All Coaches, Athletic Trainers, School Nurses, and School/Team Physicians shall complete an Interscholastic Head Injury Safety Training Program by the 2011-2012 school year.
- All school districts, charter, and non-public schools that participate in interscholastic sports will distribute
 annually this educational fact to all student athletes and obtain a signed acknowledgement from each
 parent/guardian and student-athlete.
- Each school district, charter, and non-public school shall develop a written policy describing the
 prevention and treatment of sports-related concussion and other head injuries sustained by interscholastic
 student-athletes.
- Any student-athlete who participates in an interscholastic sports program and is suspected of sustaining a
 concussion will be immediately removed from competition or practice. The student-athlete will not be
 allowed to return to competition or practice until he/she has written clearance from a physician trained in
 concussion treatment and has completed his/her district's graduated return-to-play protocol.

Ouick Facts

- Most concussions do not involve loss of consciousness
- You can sustain a concussion even if you do not hit your head
- A blow elsewhere on the body can transmit an "impulsive" force to the brain and cause a concussion

Signs of Concussions (Observed by Coach, Athletic Trainer, Parent/Guardian)

- Appears dazed or stunned
- Forgets plays or demonstrates short term memory difficulties (e.g. unsure of game, opponent)
- Exhibits difficulties with balance, coordination, concentration, and attention
- Answers questions slowly or inaccurately
- Demonstrates behavior or personality changes
- Is unable to recall events prior to or after the hit or fall

Symptoms of Concussion (Reported by Student-Athlete)

- Headache
- Nausea/vomiting
- Balance problems or dizziness
- Double vision or changes in vision

- Sensitivity to light/sound
- Feeling of sluggishness or fogginess
- Difficulty with concentration, short term memory, and/or confusion

What Should a Student-Athlete do if they think they have a concussion?

- Don't hide it. Tell your Athletic Trainer, Coach, School Nurse, or Parent/Guardian.
- Report it. Don't return to competition or practice with symptoms of a concussion or head injury. The sooner you report it, the sooner you may return-to-play.
- Take time to recover. If you have a concussion your brain needs time to heal. While your brain is
 healing you are much more likely to sustain a second concussion. Repeat concussions can cause
 permanent brain injury.

What can happen if a student-athlete continues to play with a concussion or returns to play to soon?

- Continuing to play with the signs and symptoms of a concussion leaves the student-athlete vulnerable to second impact syndrome.
- Second impact syndrome is when a student-athlete sustains a second concussion while still having symptoms from a previous concussion or head injury.
- Second impact syndrome can lead to severe impairment and even death in extreme cases.

Should there be any temporary academic accommodations made for Student-Athletes who have suffered a concussion?

- To recover cognitive rest is just as important as physical rest. Reading, texting, testing-even watching
 movies can slow down a student-athletes recovery.
- Stay home from school with minimal mental and social stimulation until all symptoms have resolved.
- Students may need to take rest breaks, spend fewer hours at school, be given extra time to complete
 assignments, as well as being offered other instructional strategies and classroom accommodations.

Student-Athletes who have sustained a concussion should complete a graduated return-to-play before they may resume competition or practice, according to the following protocol:

- Step 1: Completion of a full day of normal cognitive activities (school day, studying for tests, watching practice, interacting with peers) without reemergence of any signs or symptoms. If no return of symptoms, next day advance.
- Step 2: Light Aerobic exercise, which includes walking, swimming, and stationary cycling, keeping the
 intensity below 70% maximum heart rate. No resistance training. The objective of this step is increased
 heart rate.
- Step 3: Sport-specific exercise including skating, and/or running: no head impact activities. The objective
 of this step is to add movement.
- Step 4: Non contact training drills (e.g. passing drills). Student-athlete may initiate resistance training.
- Step 5: Following medical clearance (consultation between school health care personnel and studentathlete's physician), participation in normal training activities. The objective of this step is to restore confidence and assess functional skills by coaching and medical staff.
- Step 6: Return to play involving normal exertion or game activity.

.

www.cdc.gov/concussion/sports/in		lnjuries, please visit: www.nfhs.com	
www.ncaa.org/health-safety	www.bianj.org	www.atsnj.org	
Signature of Student-Athlete	Print Student-A	thlete's Name	Date
Signature of Parent/Guardian	Print Parent/Gu	ardian's Name	Date

Website Resources

- Sudden Death in Athletes http://tinyurl.com/m2gjmvq
- Hypertrophic Cardiomyopathy Association www.4hcm.org
- American Heart Association www.heart.org

Collaborating Agencies:

American Academy of Pediatrics New Jersey Chapter

3836 Quakerbridge Road, Suite 108 Hamilton, NJ 08619 (p) 609-842-0014 (f) 609-842-0015 www.aapnj.org

American Heart Association

1 Union Street, Suite 301 Robbinsville, NJ, 08691 (p) 609-208-0020 www.heart.org

New Jersey Department of Education

PO Box 500 Trenton, NJ 08625-0500 (p) 609-292-5935 www.state.nj.us/education/



P. O. Box 360 Trenton, NJ 08625-0360 (p) 609-292-7837 www.state.nj.us/health

Lead Author: American Academy of Pediatrics, New Jersey Chapter

Written by: Initial draft by Sushma Raman Hebbar, MD & Stephen G. Rice, MD PhD Additional Reviewers: NJ Department of Education, NJ Department of Health and Senior Services, American Heart Association/New Jersey Chapter, NJ Academy of Family Practice, Pediatric Cardiologists, New Jersey State School Nurses

Revised 2014: Nancy Curry, EdM; Christene DeWitt-Parker, MSN, CSN, RN; Lakota Kruse, MD, MPH; Susan Martz, EdM; Stephen G. Rice, MD: Jeffrey Rosenberg, MD, Louis Teichholz, MD; Perry Weinstock, MD





American Academy of Pediatrics Entrangement of ALL CHILDREN





udden death in young athletes
between the ages of 10
and 19 is very rare.
What, if anything, can be
done to prevent this kind of

What is sudden cardiac death in the young athlete?

tragedy?

Sudden cardiac death is the result of an unexpected failure of proper heart function, usually (about 60% of the time) during or immediately after exercise without trauma. Since the heart stops pumping adequately, the athlete quickly collapses, loses consciousness, and ultimately dies unless normal heart rhythm is restored using an automated external defibrillator (AED).

How common is sudden death in young athletes?

Sudden cardiac death in young athletes is very rare. About 100 such deaths are reported in the United States per year. The chance of sudden death occurring to any individual high school athlete is about one in 200,000 per year.

Sudden cardiac death is more common: in males than in females; in football and basketball than in other sports; and in African-Americans than in other races and ethnic groups.

What are the most common causes?

Research suggests that the main cause is a loss of proper heart rhythm, causing the heart to quiver instead of pumping blood to the brain and body. This is called ventricular fibrillation (ven-TRICK-you-lar fibroo-LAY-shun). The problem is usually caused by one of several cardiovascular abnormalities and electrical diseases of the heart that go unnoticed in healthy-appearing athletes.

The most common cause of sudden death in an athlete is hypertrophic cardiomyopathy (hi-per-TRO-fic CAR-dee-oh-my-OP-a-thee) also called HCM. HCM is a disease of the heart, with abnormal thickening of the heart muscle, which can cause serious heart rhythm problems and blockages to blood flow. This genetic disease runs in families and usually develops gradually over many years.

The second most likely cause is congenital (con-JEN-it-al) (i.e., present from birth) abnormalities of the coronary arteries. This means that these blood vessels are connected to the main blood vessel of the heart in an abnormal way. This differs from blockages that may occur when people get older (commonly called "coronary artery disease," which may lead to a heart attack).

Other diseases of the heart that can lead to sudden death in young people include:

- Myocarditis (my-oh-car-DIE-tis), an acute inflammation of the heart muscle (usually due to a virus).
- Dilated cardiomyopathy, an enlargement of the heart for unknown reasons.
- Long QT syndrome and other electrical abnormalities of the heart which cause abnormal fast heart rhythms that can also run in families.
- Marfan syndrome, an inherited disorder that affects heart valves, walls of major arteries, eyes and the skeleton. It is generally seen in unusually tall athletes, especially if being tall is not common in other family members.

Are there warning signs to watch for?

In more than a third of these sudden cardiac deaths, there were warning signs that were not reported or taken seriously. Warning signs are:

- Fainting, a seizure or convulsions during physical activity;
 Fainting or a seizure from emotional
 - Fainting or a seizure from emotional excitement, emotional distress or being startled;
- Dizziness or lightheadedness, especially during exertion;
- Chest pains, at rest or during exertion;
- Palpitations awareness of the heart beating unusually (skipping, irregular or extra beats) during athletics or during cool down periods after athletic participation;
- Fatigue or tiring more quickly than peers; or
- Being unable to keep up with friends due to shortness of breath (labored breathing).

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES What are the current recommendations PPE. However, these procedures may be

for screening young athletes?

New Jersey requires all school athletes to be examined by their primary care physician ("medical home") or school physician at least once per year. The New Jersey Department of Education requires use of the specific Preparticipation Physical Examination Form (PPE).

This process begins with the parents and student-athletes answering questions about symptoms during exercise (such as chest pain, dizziness, fainting, palpitations or shortness of breath); and questions about family health history.

The primary healthcare provider needs to know if any family member died suddenly during physical activity or during a seizure. They also need to know if anyone in the family under the age of 50 had an unexplained sudden death such as drowning or car accidents. This information must be provided annually for each exam because it is so essential to identify those at risk for sudden cardiac death.

The required physical exam includes measurement of blood pressure and a careful listening examination of the heart, especially for murmurs and rhythm abnormalities. If there are no warning signs reported on the health history and no abnormalities discovered on exam, no further evaluation or testing is recommended.

Are there options privately available to screen for cardiac conditions?

Technology-based screening programs including a 12-lead electrocardiogram (ECG) and echocardiogram (ECHO) are noninvasive and painless options parents may consider in addition to the required

normal screening evaluation, such as an infection of the heart muscle from a virus.

the American Academy of Pediatrics and the

expensive and are not currently advised by

addition to the expense, other limitations of

technology-based tests include the

PPE reveals an indication for these tests. In

American College of Cardiology unless the

possibility of "false positives" which leads to

parent or guardian as well as unnecessary

restriction from athletic participation.

unnecessary stress for the student and

This is why screening evaluations and a review of the family health history need to be performed on a yearly basis by the athlete's primary healthcare provider. With proper screening and evaluation, most cases can be identified and prevented.

Why have an AED on site during sporting

The only effective treatment for ventricular fibrillation is immediate use of an automated external defibrillator (AED). An AED can restore the heart back into a normal rhythm. An AED is also life-saving for ventricular fibrillation caused by a blow to the chest over the heart (commotio cordis).

http://www.hhs.gov/familyhistory/index.html.

History Initiative available at

When should a student athlete see a

heart specialist?

options under the Surgeon General's Family

and Human Services offers risk assessment

The United States Department of Health

NJ.S.A. 18A:40-41a through c, known as "Janet's Law," requires that at any schoolsponsored athletic event or team practice in New Jersey public and nonpublic schools including any of grades K through 12, the following must be available:

 An AED in an unlocked location on school property within a reasonable proximity to the athletic field or gymnasium; and

electrocardiogram (ECG), which is a graph of

a more thorough evaluation, including an

recommended. This specialist will perform

heart specialist, a pediatric cardiologist, is

If the primary healthcare provider or schoo

physician has concerns, a referral to a child

echocardiogram, which is an ultrasound test

the electrical activity of the heart. An

to allow for direct visualization of the heart

structure, will likely also be done. The

 A team coach, licensed athletic trainer, or other designated staff member if there is no coach or licensed athletic trainer present, certified in cardiopulmonary resuscitation (CPR) and the use of the AED; or

specialist may also order a treadmill exercise

recording of the heart rhythm. None of the

testing is invasive or uncomfortable.

test and a monitor to enable a longer

A State-certified emergency services provider or other certified first responder.

The American Academy of Pediatrics recommends the AED should be placed in central location that is accessible and ideally no more than a 1 to $1^{1/2}$ minute walk from any location and that a call is made to activate 911 emergency system while the AED is being retrieved.

A proper evaluation should find most, but not

Can sudden cardiac death be prevented

just through proper screening?

all, conditions that would cause sudden death

in the athlete. This is because some diseases

are difficult to uncover and may only develop

ater in life. Others can develop following a

State of New Jersey DEPARTMENT OF EDUCATION

Sudden Cardiac Death Pamphlet Sign-Off Sheet

Name of School District:
Name of Local School:
I/We acknowledge that we received and reviewed the Sudden Cardiac Death in Young Athletes pamphlet.
Student Signature:
Parent or Guardian
Signature:
Date: