

Ocean City Public Schools Pre-Participation Forms



Student Name	Gender	Grade	Sport
HISTORY REVIEWED AND S	STUDENT EXAMINED	BY:	Physician's/Provider's Stamp:
☐ Primary Care Provider			
☐ School Physician Provider			
☐ License Type:			
□ MD/DO			
\Box APN			
\Box PA			
PHYSICIAN'S/PROVIDER'S SIGNAT	TURE:		
Today's Date:	Date of Exam:		
	DIAN of the child listed aboves Registration and agree to all	eancitys s Regist e, I con guideli	ration" and follow instructions npleted the online portion of nes and policies mentioned.
☐ <u>Use and Misuse of Opioid Drug</u>	gs Fact Sheet		
☐ Opioid Education Video Procee	<u>dure¹</u>		
☐ Sports-Related Concussion and	Head Injury Fact Sheet		
☐ Sudden Cardiac Death in Youn	ng Athletes Pamphlet		
Parent/Guardian Name:			
Parent/Guardian Signature:			
Date:Please return this form along with the by the first day of try-outs. Any studer - NO EXCEPTIONS!	other sign-off sheets listed a nt that does not return this f 1	above to form wi	o the above to the Health Office ll not be allowed to participate –

ATTENTION PARENT/GUARDIAN: The preparticipation physical examination (page 3) must be completed by a health care provider who has completed the Student-Athlete Cardiac Assessment Professional Development Module.

■ PREPARTICIPATION PHYSICAL EVALUATION

HISTORY FORM

me	Date of birth				
	School Sport(s)				
			edicines and supplements (herbal and nutritional) that you are currently	tokina	_
nedictines and Allergies: Please list all of the prescription and over	-tne-co	unter m	edicines and supplements (nerbal and nutritional) that you are currently	taking	
					Ξ
					_
00 you have any allergies?	ntify spe	ecific all	lergy below. □ Food □ Stinging Insects		_
plain "Yes" answers below. Circle questions you don't know the an	_		MEDICAL CHECKTONS	W	Т
ENERAL QUESTIONS	Yes	No	MEDICAL QUESTIONS 26. Do you cough, wheeze, or have difficulty breathing during or	Yes	ł
1. Has a doctor ever denied or restricted your participation in sports for any reason?			after exercise?		ļ
2. Do you have any ongoing medical conditions? If so, please identify			27. Have you ever used an inhaler or taken asthma medicine?		ļ
below: Asthma Anemia Diabetes Infections Other:			28. Is there anyone in your family who has asthma?	_	ł
. Have you ever spent the night in the hospital?			29. Were you born without or are you missing a kidney, an eye, a testicle (males), your spleen, or any other organ?		
. Have you ever had surgery?			30. Do you have groin pain or a painful bulge or hernia in the groin area?		Ι
EART HEALTH QUESTIONS ABOUT YOU	Yes	No	31. Have you had infectious mononucleosis (mono) within the last month?		ſ
i. Have you ever passed out or nearly passed out DURING or AFTER exercise?			32. Do you have any rashes, pressure sores, or other skin problems?		ĺ
			33. Have you had a herpes or MRSA skin infection?		ļ
6. Have you ever had discomfort, pain, tightness, or pressure in your chest during exercise?			34. Have you ever had a head injury or concussion?		ļ
. Does your heart ever race or skip beats (irregular beats) during exercise?			35. Have you ever had a hit or blow to the head that caused confusion, prolonged headache, or memory problems?		l
3. Has a doctor ever told you that you have any heart problems? If so,			36. Do you have a history of seizure disorder?		t
check all that apply: High blood pressure			37. Do you have headaches with exercise?		t
☐ High blood pressure ☐ A heart murmur ☐ High cholesterol ☐ A heart infection			38. Have you ever had numbness, tingling, or weakness in your arms or		t
☐ Kawasaki disease Other:			legs after being hit or falling?		ļ
Has a doctor ever ordered a test for your heart? (For example, ECG/EKG, echocardiogram)			39. Have you ever been unable to move your arms or legs after being hit or falling?		
Do you get lightheaded or feel more short of breath than expected			40. Have you ever become ill while exercising in the heat?		l
during exercise?			41. Do you get frequent muscle cramps when exercising?		ļ
Have you ever had an unexplained seizure?			42. Do you or someone in your family have sickle cell trait or disease?	_	Ļ
2. Do you get more tired or short of breath more quickly than your friends during exercise?			43. Have you had any problems with your eyes or vision?		ł
EART HEALTH QUESTIONS ABOUT YOUR FAMILY	Yes	No	44. Have you had any eye injuries?		ł
3. Has any family member or relative died of heart problems or had an			45. Do you wear glasses or contact lenses?		ł
unexpected or unexplained sudden death before age 50 (including drowning, unexplained car accident, or sudden infant death syndrome)?			46. Do you wear protective eyewear, such as goggles or a face shield? 47. Do you worry about your weight?		ł
4. Does anyone in your family have hypertrophic cardiomyopathy, Marfan			48. Are you trying to or has anyone recommended that you gain or		Γ
syndrome, arrhythmogenic right ventricular cardiomyopathy, long QT			lose weight?		ļ
syndrome, short QT syndrome, Brugada syndrome, or catecholaminergic polymorphic ventricular tachycardia?			49. Are you on a special diet or do you avoid certain types of foods?		+
5. Does anyone in your family have a heart problem, pacemaker, or			50. Have you ever had an eating disorder?	 	+
implanted defibrillator?			51. Do you have any concerns that you would like to discuss with a doctor? FEMALES ONLY		+
6. Has anyone in your family had unexplained fainting, unexplained seizures, or near drowning?			52. Have you ever had a menstrual period?		+
ONE AND JOINT QUESTIONS	Yes	No	53. How old were you when you had your first menstrual period?		L
7. Have you ever had an injury to a bone, muscle, ligament, or tendon	105		54. How many periods have you had in the last 12 months?		-
that caused you to miss a practice or a game?			Explain "yes" answers here		-
8. Have you ever had any broken or fractured bones or dislocated joints?					
9. Have you ever had an injury that required x-rays, MRI, CT scan,					
injections, therapy, a brace, a cast, or crutches?					
Have you ever had a stress fracture?					
 Have you ever been told that you have or have you had an x-ray for neck instability or atlantoaxial instability? (Down syndrome or dwarfism) 					_
Do you regularly use a brace, orthotics, or other assistive device?					_
Do you have a bone, muscle, or joint injury that bothers you?					_
Do any of your joints become painful, swollen, feel warm, or look red?					-
5. Do you have any history of juvenile arthritis or connective tissue disease?	I	1			

Signature of athlete Signature of parent/guardian Date

© 2010 American Academy of Earthy Physicians American Academy of Badistrics American College of Sports Madicine American Medical Society for Sports Madicine American Orthogoadis

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

■ PREPARTICIPATION PHYSICAL EVALUATION

THE ATHLETE WITH SPECIAL NEEDS: SUPPLEMENTAL HISTORY FORM

Date	of Exam					
Name				Dat	e of birth	
		O-refe	Cohool			
Sex _	Age	Grade	School	Sport(s)		
1. To	ype of disability					
-	ate of disability					
-	lassification (if available)					
		ease, accident/trauma, other	2			
-	st the sports you are intere)			
5. LI	as the sports you are intere	ested in playing			Yes	No
6 D	o vou requilarly use a brace	e, assistive device, or prosthe	atic?		165	140
-		e or assistive device for spor				
-		ssure sores, or any other ski				
-			ii problems:			
	o you have a rearing loss? o you have a visual impair	Do you use a hearing aid?				
-		ces for bowel or bladder fund	rtion?			
-	o you use any special devi o you have burning or disc		cuone			
-						
-	ave you had autonomic dys		athermia) as and extend formathermia) floor	-0		
-			rthermia) or cold-related (hypothermia) illnes	8?		
-	o you have muscle spastic	-				
16. D	o you have frequent seizur	es that cannot be controlled	by medication?			
Explain	n "yes" answers here					
_						
_						
Please	indicate if you have ever	r had any of the following.				
	maioate ii jou nave ever	nou any or are renowing.				
					Voc	Mo
Atlant	navial inetahility				Yes	No
-	naxial instability	inetahilitu			Yes	No
X-ray	evaluation for atlantoaxial				Yes	No
X-ray Disloc	evaluation for atlantoaxial ated joints (more than one				Yes	No
X-ray Disloc Easy t	evaluation for atlantoaxial sated joints (more than one bleeding				Yes	No
X-ray Disloc Easy t Enlarg	evaluation for atlantoaxial ated joints (more than one bleeding ged spleen				Yes	No
X-ray Disloc Easy t Enlarg Hepat	evaluation for atlantoaxial ated joints (more than one) bleeding ged spleen itis				Yes	No
X-ray Disloc Easy t Enlarg Hepat Osteo	evaluation for atlantoaxial ated joints (more than one) bleeding ged spleen itits penia or osteoporosis				Yes	No
X-ray Disloc Easy t Enlarg Hepat Osteo Difficu	evaluation for atlantoaxial ated joints (more than one) bleeding ged spleen itits penia or osteoporosis ulty controlling bowel				Yes	No
X-ray Disloc Easy I Enlarg Hepat Osteo Difficu	evaluation for atlantoaxial ated joints (more than one bleeding ged spleen itis penia or osteoporosis ulty controlling bowel ulty controlling bladder)			Yes	No
X-ray Disloc Easy t Enlarg Hepat Osteo Difficu Numb	evaluation for atlantoaxial ated joints (more than one bleeding ged spleen itis penia or osteoporosis ulty controlling bowel ulty controlling bladder ness or tingling in arms or	hands			Yes	No
X-ray Disloc Easy t Enlarg Hepat Osteo Difficu Numb	evaluation for attantoaxial sated joints (more than one bleeding get spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder iness or tingling in arms or iness or tingling in legs or f	hands			Yes	No
X-ray Disloc Easy I Enlarg Hepat Osteo Difficu Numb Weak	evaluation for atlantoaxial sated joints (more than one bleeding ged spleen jed spleen j	hands			Yes	No
X-ray Disloc Easy I Enlarg Hepat Osteo Difficu Numb Numb Weak	evaluation for attantoaxial sated joints (more than one bleeding ged spleen ittis penia or osteoporosis uity controlling bowel uity controlling bladder mess or tingling in arms or mess or tingling in legs or f mess in arms or hands mess in legs or feet	hands			Yes	No
X-ray Disloc Easy t Enlarg Hepat Osteo Difficu Numb Numb Weak Weak	evaluation for attantoaxial sated joints (more than one bleeding ged spleen itis penia or osteoporosis alty controlling bowel alty controlling bladder mess or tingling in arms or mess or tingling in legs or finess in arms or hands mess in legs or feet at change in coordination	hands			Yes	No
X-ray Disloc Easy I Enlarg Hepat Osteo Difficu Numb Weak Weak Recen	evaluation for attantoaxial sated joints (more than one bleeding ged spleen itis penia or osteoporosis sity controlling bowel sity controlling bladder mess or tingling in arms or mess or tingling in legs or finess in arms or hands mess in legs or feet to thange in coordination at change in ability to walk	hands			Yes	No
X-ray Disloc Easy I Enlarg Hepat Osteo Diffica Numb Weak Weak Recen Recen Spina	evaluation for attantoaxial sated joints (more than one bleeding ged spleen itis penia or osteoporosis uity controlling bowel uity controlling bladder mess or tingling in arms or mess in legs or feet at change in coordination at change in ability to walk bifida	hands			Yes	No
X-ray Disloc Easy I Enlarg Hepat Osteo Diffica Numb Weak Weak Recen Recen Spina	evaluation for attantoaxial sated joints (more than one bleeding ged spleen itis penia or osteoporosis sity controlling bowel sity controlling bladder mess or tingling in arms or mess or tingling in legs or finess in arms or hands mess in legs or feet to thange in coordination at change in ability to walk	hands			Yes	No
X-ray Disloce Easy I Enlarge Hepat Osteo Difficu Numb Weako Weako Recer Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ged spleen itis penia or osteoporosis uity controlling bowel uity controlling bladder mess or tingling in arms or mess in legs or feet at change in coordination at change in ability to walk bifida	hands			Yes	No
X-ray Disloct Easy Mental Enlarge Hepat Osteo Difficution Numb Weako Weako Recent Recent Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ded spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder siness or tingling in arms or siness or tingling in legs or tiness in arms or hands siness in legs or feet at change in coordination at change in ability to walk bifida allergy	hands			Yes	No
X-ray Disloct Easy Mental Enlarge Hepat Osteo Difficution Numb Weako Weako Recent Recent Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ded spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder siness or tingling in arms or siness or tingling in legs or tiness in arms or hands siness in legs or feet at change in coordination at change in ability to walk bifida allergy	hands			Yes	No
X-ray Disloct Easy Mental Enlarge Hepat Osteo Difficution Numb Weako Weako Recent Recent Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ded spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder siness or tingling in arms or siness or tingling in legs or tiness in arms or hands siness in legs or feet at change in coordination at change in ability to walk bifida allergy	hands			Yes	No
X-ray Disloce Easy I Enlarge Hepat Osteo Difficu Numb Weako Weako Recer Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ded spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder siness or tingling in arms or siness or tingling in legs or tiness in arms or hands siness in legs or feet at change in coordination at change in ability to walk bifida allergy	hands			Yes	No
X-ray Disloct Easy Mental Enlarge Hepat Osteo Difficution Numb Weako Weako Recent Recent Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ded spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder siness or tingling in arms or siness or tingling in legs or tiness in arms or hands siness in legs or feet at change in coordination at change in ability to walk bifida allergy	hands			Yes	No
X-ray Disloce Easy I Enlarge Hepat Osteo Difficu Numb Weako Weako Recer Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ded spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder siness or tingling in arms or siness or tingling in legs or tiness in arms or hands siness in legs or feet at change in coordination at change in ability to walk bifida allergy	hands			Yes	No
X-ray Disloct Easy Mental Enlarge Hepat Osteo Difficution Numb Weako Weako Recent Recent Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ded spleen sitis penia or osteoporosis sity controlling bowel sity controlling bladder siness or tingling in arms or siness or tingling in legs or tiness in arms or hands siness in legs or feet at change in coordination at change in ability to walk bifida allergy	hands			Yes	No
X-ray Disloc Easy I Enlarg Hepat Osteo Difficu Numb Weak Weak Recen Spina Latex	evaluation for attantoaxial sated joints (more than one bleeding ged spleen it is penia or osteoporosis alty controlling bowel alty controlling bladder mess or tingling in arms or mess or tingling in legs or finess in arms or hands mess in legs or feet at change in coordination at change in ability to walk bifida allergy	hands feet	vers to the above questions are complete	and correct.	Yes	No
X-ray Disloc Easy to Enlarge Hepat Osteo Difficution Difficution Numb Weak Weak Recent Recent Spina Latex Explain	evaluation for attantoaxial sated joints (more than one bleeding ged spleen it is penia or osteoporosis alty controlling bowel alty controlling bladder mess or tingling in arms or mess or tingling in legs or finess in arms or hands mess in legs or feet at change in coordination at change in ability to walk bifida allergy	hands feet	ers to the above questions are complete a	and correct.	Yes Date	No

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71

NOTE: The preparticiaption physical examination must be conducted by a health care provider who 1) is a licensed physician, advanced practice nurse, or physician assistant; and 2) completed the Student-Athlete Cardiac Assessment Professional Development Module.

Date of birth

■ PREPARTICIPATION PHYSICAL EVALUATION

PHYSICAL EXAMINATION FORM

Name

PHYSICIAN REMINDERS

1. Consider additional questions on more sensitive issues Do you feel stressed out or under a lot of pressure? Do you ever feel sad, hopeless, depressed, or anxious? Do you feel safe at your home or residence? Have you ever tried cigarettes, chewing tobacco, snuff, or dip? . During the past 30 days, did you use chewing tobacco, snuff, or dip? * Do you drink alcohol or use any other drugs? * Have you ever taken anabolic steroids or used any other performance supplement? Have you ever taken any supplements to help you gain or lose weight or improve your performance?
 Do you wear a seat belt, use a helmet, and use condoms? 2. Consider reviewing questions on cardiovascular symptoms (questions 5-14). **EXAMINATION** ■ Male ■ Female Height BP Pulse Vision R 20/ 1.20/ Corrected Y \square N MEDICAL ABNORMAL FINDINGS NORMAL Marfan stigmata (kyphoscoliosis, high-arched palate, pectus excavatum, arachnodactyly, arm span > height, hyperlaxity, myopia, MVP, aortic insufficiency) Eyes/ears/nose/throat Pupils equal Hearing Lymph nodes · Murmurs (auscultation standing, supine, +/- Valsalva) · Location of point of maximal impulse (PMI) Pulses · Simultaneous femoral and radial pulses Lungs Abdomen Genitourinary (males only)^b Skin . HSV, lesions suggestive of MRSA, tinea corporis Neurologic MUSCULOSKELETAL Neck Back Shoulder/arm Elbow/forearm Wrist/hand/fingers Hip/thigh Knee Leg/ankle Foot/toes Functional · Duck-walk, single leg hop *Consider ECG, echocardiogram, and referral to cardiology for abnormal cardiac history or exam. *Consider GU exam if in private setting, Having third party present is recommended. *Consider cognitive eval ation or baseline neuropsychiatric testing if a history of significant concussion. Cleared for all sports without restriction ☐ Cleared for all sports without restriction with recommendations for further evaluation or treatment for □ Not cleared □ Pending further evaluation For any sports ☐ For certain sports _____ Recommendations I have examined the above-named student and completed the preparticipation physical evaluation. The athlete does not present apparent clinical contraindications to practice and participate in the sport(s) as outlined above. A copy of the physical exam is on record in my office and can be made available to the school at the request of the parents. If conditions arise after the athlete has been cleared for participation, a physician may rescind the clearance until the problem is resolved and the potential consequences are completely explained to the athlete (and parents/guardians). Name of physician, advanced practice nurse (APN), physician assistant (PA) (print/type)_ Date of exam ___

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

Signature of physician, APN, PA

■ PREPARTICIPATION PHYSICAL EVALUATION CLEARANCE FORM

Name	Sex M F Age Date of birth
☐ Cleared for all sports without restriction	
Cleared for all sports without restriction with recommendations for further evaluations.	luation or treatment for
□ Not cleared	
 Pending further evaluation 	
☐ For any sports	
☐ For certain sports	
Reason	
Recommendations	
EMERGENCY INFORMATION	
Allergies	
Other Information	
Other information	
	aguasi suugasuu
HCP OFFICE STAMP	SCHOOL PHYSICIAN:
	Reviewed on(Date)
	Approved Not Approved
	Signature:
	Signature:
clinical contraindications to practice and participate in the sport(s) and can be made available to the school at the request of the paren	articipation physical evaluation. The athlete does not present apparent as outlined above. A copy of the physical exam is on record in my office ts. If conditions arise after the athlete has been cleared for participation, ed and the potential consequences are completely explained to the athlete
Name of physician, advanced practice nurse (APN), physician assistant (PA)	Date
Address	Phone
Signature of physician, APN, PA	
Completed Cardiac Assessment Professional Development Module	
Date Signature	

© 2010 American Academy of Family Physicians, American Academy of Pediatrics, American College of Sports Medicine, American Medical Society for Sports Medicine, American Orthopaedic Society for Sports Medicine, and American Osteopathic Academy of Sports Medicine. Permission is granted to reprint for noncommercial, educational purposes with acknowledgment.

New Jersey Department of Education 2014; Pursuant to P.L.2013, c.71



OPIOID USE AND MISUSE EDUCATIONAL FACT SHEET

Keeping Student-Athletes Safe

School athletics can serve an integral role in students' development. In addition to providing healthy forms of exercise, school athletics foster friendships and camaraderie, promote sportsmanship and fair play, and instill the value of competition.

Unfortunately, sports activities may also lead to injury and, in rare cases, result in pain that is severe or long-lasting enough to require a prescription opioid painkiller. It is important to understand that overdoses from opioids are on the rise and are killing Americans of all ages and backgrounds. Families and communities across the country are coping with the health, emotional and economic effects of this epidemic.²

This educational fact sheet, created by the New Jersey Department of Education as required by state law (*N.J.S.A.* 18A:40-41.10), provides information concerning the use and misuse of opioid drugs in the event that a health care provider prescribes a student-athlete or cheerleader an opioid for a sports-related injury. Student-athletes and cheerleaders participating in an interscholastic sports program (and their parent or guardian, if the student is under age 18) must provide their school district written acknowledgment of their receipt of this fact sheet.

How Do Athletes Obtain Opioids?

In some cases, student-athletes are prescribed these medications. According to research, about a third of young people studied obtained pills from their own previous prescriptions (i.e., an unfinished prescription used outside of a physician's supervision), and 83 percent of adolescents had unsupervised access to their prescription medications. It is important for parents to understand the possible hazard of having unsecured prescription medications in their households. Parents should also understand the importance of proper storage and disposal of medications, even if they believe their child would not engage in non-medical use or diversion of prescription medications.

What Are Signs of Opioid Use?

According to the National Council on Alcoholism and Drug Dependence, 12 percent of male athletes and 8 percent of female athletes had used prescription opioids in the 12-month period studied.³ In the early stages of abuse, the athlete may exhibit unprovoked nausea and/or vomiting. However, as he or she develops a tolerance to the drug, those signs will diminish. Constipation is not uncommon, but may not be reported. One of the most significant indications of a possible opioid addiction is an athlete's decrease in academic or athletic performance, or a lack of interest in his or her sport. If these warning signs are noticed, best practices call for the student to be referred to the appropriate professional for screening, ⁴ such as provided through an evidence-based practice to identify problematic use, abuse and dependence on illicit drugs (e.g., Screening, Brief Intervention, and Referral to Treatment (SBIRT)) offered through the New Jersey Department of Health.

What Are Some Ways Opioid Use and Misuse Can Be Prevented?

According to the New Jersey State Interscholastic Athletic Association (NJSIAA) Sports Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcotic painkillers."

Medical Advisory Committee chair, John P. Kripsak, D.O., "Studies indicate that about 80 percent of heroin users started out by abusing narcatic painkillers."

According to NJSIAA Sports

The Sports Medical Advisory Committee, which includes representatives of NJSIAA member schools as well as experts in the field of healthcare and medicine, recommends the following:

- The pain from most sports-related injuries can be managed with non-narcotic medications such as acetaminophen, nonsteroidal anti-inflammatory medications like ibuprofen, naproxen or aspirin. Read the label carefully and always take the recommended dose, or follow your doctor's instructions. More is not necessarily better when taking an over-the-counter (OTC) pain medication, and it can lead to dangerous side effects.
- Ice therapy can be utilized appropriately as an anesthetic.
- Always discuss with your physician exactly what is being prescribed for pain and request to avoid narcotics.
- In extreme cases, such as severe trauma or post-surgical pain, opioid pain medication should not be prescribed for more than five days at a time;
- Parents or guardians should always control the dispensing of pain medications and keep them in a safe, non-accessible location; and
- Unused medications should be disposed of immediately upon cessation of use. Ask your pharmacist about drop-off locations
 or home disposal kits like Deterra or Medsaway.






STATE OF NEW JERSEY DEPARTMENT OF HEALTH

NISIAA SPORTS MEDICAL ADVISORY COMMITTEE





Even With Proper Training and Prevention, Sports Injuries May Occur

There are two kinds of sports injuries. Acute injuries happen suddenly, such as a sprained ankle or strained back. Chronic injuries may happen after someone plays a sport or exercises over a long period of time, even when applying overuse-preventative techniques.5

Athletes should be encouraged to speak up about injuries, coaches should be supported in injury-prevention decisions, and parents and young athletes are encouraged to become better educated about sports safety.6

What Are Some Ways to Reduce the Risk of Injury?

Half of all sports medicine injuries in children and teens are from overuse. An overuse injury is damage to a bone, muscle, ligament, or tendon caused by repetitive stress without allowing time for the body to heal. Children and teens are at increased risk for overuse injuries because growing bones are less resilient to stress. Also, young athletes may not know that certain symptoms are signs of overuse.

The best way to deal with sports injuries is to keep them from happening in the first place. Here are some recommendations to consider:



PREPARE Obtain the preparticipation physical evaluation prior to participation on a school-sponsored interscholastic or intramural athletic team or squad.



CONDITIONING Maintain a good fitness level during the season and offseason. Also important are proper warm-up and cooldown



PLAY SMART Try a variety of sports and consider specializing in one sport before late adolescence to help avoid overuse injuries.



ADEQUATE HYDRATION Keep the body hydrated to help the heart more easily pump blood to muscles, which helps muscles work efficiently.



TRAINING Increase weekly training time, mileage or repetitions no more than 10 percent per week. For example, if running 10 miles one week, increase to 11 miles the following week. Athletes should also cross-train and perform sport-specific drills in different ways, such as running in a swimming pool instead of only running on the road.



REST UP Take at least one day off per week from organized activity to recover physically and mentally. Athletes should take a combined three months off per year from a specific sport (may be divided throughout the year in one-month increments). Athletes may remain physically active during rest periods through alternative low-stress activities such as stretching, yoga or walking.



PROPER EQUIPMENT Wear appropriate and properly fitted protective equipment such as pads (neck, shoulder, elbow, chest, knee, and shin), helmets, mouthpieces, face guards, protective cups, and eyewear. Do not assume that protective gear will prevent all injuries while performing more dangerous or risky activities.

Resources for Parents and Students on Preventing Substance Misuse and Abuse

The following list provides some examples of resources:

National Council on Alcoholism and Drug Dependence - NJ promotes addiction treatment and recovery.

New Jersey Department of Health, Division of Mental Health and Addiction Services is committed to providing consumers and families with a wellness and recovery-oriented model of care.

New Jersey Prevention Network includes a parent's quiz on the effects of opioids.

Operation Prevention Parent Toolkit is designed to help parents learn more about the opioid epidemic, recognize warning signs, and open lines of communication with their children and those in the community.

Parent to Parent NJ is a grassroots coalition for families and children struggling with alcohol and drug addiction.

Partnership for a Drug Free New Jersey is New Jersey's anti-drug alliance created to localize and strengthen drug-prevention media efforts to prevent unlawful drug use, especially among young people.

The Science of Addiction: The Stories of Teens shares common misconceptions about opioids through the voices of teens.

Youth IMPACTing NJ is made up of youth representatives from coalitions across the state of New Jersey who have been impacting their communities and peers by spreading the word about the dangers of underage drinking, marijuana use, and other substance misuse.

- References 1 Massachusetts Technical Assistance Partnership
 - for Prevention ² Centers for Disease Control and Prevention
 - ³ New Jersey State Interscholastic Athletic
- Association (NJSIAA) Sports Medical Advisory Committee (SMAC)
- 4 Athletic Management, David Csillan, athletic trainer, Ewing High School, NJSIAA SMAC
- 5 National Institute of Arthritis and Musculoskeletal
- and Skin Diseases
- # USA TODAY
- 7 American Academy of Pediatrics

An online version of this fact sheet is available on the New Jersey Department of Education's Alcohol, Tobacco, and Other Drug Use webpage. Updated Jan. 30, 2018.

OCEAN CITY SCHOOL DISTRICT

501 Atlantic Avenue, Suite 1 Ocean City, New Jersey 08226 _ 3891

Phone: (609) 399-5150 www.oceancityschools.org



Use and Misuse of Opioid Drugs Fact Sheet Student-Athlete and Parent/Guardian Sign-Off

In accordance with N.J.S.A. 18A:40-41.10, public school districts, approved private schools for students with disabilities, and nonpublic schools participating in an interscholastic sports program must distribute the <u>Opioid Use and Misuse Educational Fact Sheet</u> to all student-athletes and cheerleaders. In addition, schools and districts must obtain a signed acknowledgement of receipt of the fact sheet from each student-athlete and cheerleader, and for students under age 18, the parent or guardian must also sign.

This sign-off sheet is due to the Health Office prior to the first official practice session of every student-athlete or cheerleader. This acknowledgement is required before <u>each</u> season in which the student-athlete or cheerleader will be participating.

Name of School: Ocean City High School & Intermediate School
Name of School District (if applicable): Ocean City School District
I/We acknowledge that we received and reviewed the online version of the Educational Fact Sheet on the Use and Misuse of Opioid Drugs while completing the online sign-up form for sports within the Ocean City School District.
Student Name:
Student Signature:
Parent/Guardian Signature:
Date:

NEW JERSEY STATE INTERSCHOLASTIC ATHLETIC ASSOCIATION

1161 Route 130 North, Robbinsville, NJ 08691 Phone 609-259-2776 ~ Fax 609-259-3047

Memorandum

To: All Athletic Directors of Member Schools

From: Tony Maselli, Assistant Director

Date: June 2019

Re: Opioid Education Video Procedure

To All Athletic Directors:

Acting to address the increased risk of opioid abuse among high school athletes, the Office of the New Jersey Coordinator for Addiction Responses and Enforcement Strategies (NJCARES) and the New Jersey State Interscholastic Athletic Association (NJSIAA) announced on February 19, 2019, a new partnership to educate student athletes and their parents/guardians on addiction risks associated with sports injuries and opioid use.

This educational initiative, spearheaded by Attorney General Gurbir Grewal and approved by the Executive Committee of the NJSIAA, is a collaborative effort to use video programming to raise awareness among high school athletes that they face a higher risk of becoming addicted to prescription pain medication than their fellow students who do not play sports.

Beginning with the 2019 fall season, we are making available to all student athletes and their parents/guardians, an educational video about the risks of opioid use as it relates to student athletes. The video will be available on August 1, 2019 and can be found on the NJSIAA website under "Athlete Wellness" which is located under the "Health & Safety tab. We are strongly encouraging student athletes and parents/guardians to watch the video as soon as it becomes available. An acknowledgement that students and their parents/guardians have watched the video will be required starting with the 2019-2020 winter season.

All member schools are asked to add to their current athletic consent forms the sign-off listed below. The sign-off acknowledgment is an NJSIAA mandate; student athletes are required to view the video only once per school year prior to the first official practice of the season in their respective sport, but the signed acknowledgment is required for each sport a student participates in. Athletes that are 18 years or older do not need the parents/guardians to watch the video.

Opioid Video is located at: https://youtu.be/3Rz6rkwpAx8

NJSIAA OPIOID POLICY ACKNOWLEDGEMENT

We have viewed the NJ CARES educational video on the risks of opioid use for high school athletes. We understand the NJSIAA policy that requires students, and their parents(s)/guardian(s) if a student is under the age of 18, to view this video and sign this acknowledgement.

Student's Signature:	Date:		
<u> </u>			
Parent/Guardian Signature:	Date:		

Sports-Related Concussion and Head Injury Fact Sheet and Parent/Guardian Acknowledgement Form

A concussion is a brain injury that can be caused by a blow to the head or body that disrupts normal functioning of the brain. Concussions are a type of Traumatic Brain Injury (TBI), which can range from mild to severe and can disrupt the way the brain normally functions. Concussions can cause significant and sustained neuropsychological impairment affecting problem solving, planning, memory, attention, concentration, and behavior.

The Centers for Disease Control and Prevention estimates that 300,000 concussions are sustained during sports related activities nationwide, and more than 62,000 concussions are sustained each year in high school contact sports. Second-impact syndrome occurs when a person sustains a second concussion while still experiencing symptoms of a previous concussion. It can lead to severe impairment and even death of the victim.

Legislation (P.L. 2010, Chapter 94) signed on December 7, 2010, mandated measures to be taken in order to ensure the safety of K-12 student-athletes involved in interscholastic sports in New Jersey. It is imperative that athletes, coaches, and parent/guardians are educated about the nature and treatment of sports related concussions and other head injuries. The legislation states that:

- All Coaches, Athletic Trainers, School Nurses, and School/Team Physicians shall complete an Interscholastic Head Injury Safety Training Program by the 2011-2012 school year.
- All school districts, charter, and non-public schools that participate in interscholastic sports will distribute
 annually this educational fact to all student athletes and obtain a signed acknowledgement from each
 parent/guardian and student-athlete.
- Each school district, charter, and non-public school shall develop a written policy describing the
 prevention and treatment of sports-related concussion and other head injuries sustained by interscholastic
 student-athletes.
- Any student-athlete who participates in an interscholastic sports program and is suspected of sustaining a
 concussion will be immediately removed from competition or practice. The student-athlete will not be
 allowed to return to competition or practice until he/she has written clearance from a physician trained in
 concussion treatment and has completed his/her district's graduated return-to-play protocol.

Ouick Facts

- · Most concussions do not involve loss of consciousness
- You can sustain a concussion even if you do not hit your head
- A blow elsewhere on the body can transmit an "impulsive" force to the brain and cause a concussion

Signs of Concussions (Observed by Coach, Athletic Trainer, Parent/Guardian)

- Appears dazed or stunned
- · Forgets plays or demonstrates short term memory difficulties (e.g. unsure of game, opponent)
- · Exhibits difficulties with balance, coordination, concentration, and attention
- Answers questions slowly or inaccurately
- Demonstrates behavior or personality changes
- · Is unable to recall events prior to or after the hit or fall

Symptoms of Concussion (Reported by Student-Athlete)

- Headache
- Nausea/vomiting
- Balance problems or dizziness
- Double vision or changes in vision

- Sensitivity to light/sound
- Feeling of sluggishness or fogginess
- Difficulty with concentration, short term memory, and/or confusion

What Should a Student-Athlete do if they think they have a concussion?

- . Don't hide it. Tell your Athletic Trainer, Coach, School Nurse, or Parent/Guardian.
- Report it. Don't return to competition or practice with symptoms of a concussion or head injury. The sooner you report it, the sooner you may return-to-play.
- Take time to recover. If you have a concussion your brain needs time to heal. While your brain is healing
 you are much more likely to sustain a second concussion. Repeat concussions can cause permanent brain
 injury.

What can happen if a student-athlete continues to play with a concussion or returns to play to soon?

- Continuing to play with the signs and symptoms of a concussion leaves the student-athlete vulnerable to second impact syndrome.
- Second impact syndrome is when a student-athlete sustains a second concussion while still having symptoms from a previous concussion or head injury.
- Second impact syndrome can lead to severe impairment and even death in extreme cases.

Should there be any temporary academic accommodations made for Student-Athletes who have suffered a concussion?

- To recover cognitive rest is just as important as physical rest. Reading, texting, testing-even watching
 movies can slow down a student-athletes recovery.
- Stay home from school with minimal mental and social stimulation until all symptoms have resolved.
- Students may need to take rest breaks, spend fewer hours at school, be given extra time to complete
 assignments, as well as being offered other instructional strategies and classroom accommodations.

Student-Athletes who have sustained a concussion should complete a graduated return-to-play before they may resume competition or practice, according to the following protocol:

- Step 1: Completion of a full day of normal cognitive activities (school day, studying for tests, watching
 practice, interacting with peers) without reemergence of any signs or symptoms. If no return of symptoms,
 next day advance.
- Step 2: Light Aerobic exercise, which includes walking, swimming, and stationary cycling, keeping the
 intensity below 70% maximum heart rate. No resistance training. The objective of this step is increased
 heart rate.
- Step 3: Sport-specific exercise including skating, and/or running: no head impact activities. The objective
 of this step is to add movement.
- Step 4: Non-contact training drills (e.g. passing drills). Student-athlete may initiate resistance training.
- Step 5: Following medical clearance (consultation between school health care personnel and studentathlete's physician), participation in normal training activities. The objective of this step is to restore confidence and assess functional skills by coaching and medical staff.
- Step 6: Return to play involving normal exertion or game activity.

For further information on Sports-Related Concussions and other Head Injuries, please visit:

- CDC Heads Up
- Keeping Heads Healthy
- National Federation of State High School Associations
- Athletic Trainers' Society of New Jersey

Signature of Student-Athlete	Print Student-Athlete's Name	Date		
Signature of Parent/Guardian	Print Parent/Guardian's Name	Date		

Website Resources

- http://tinyurl.com/m2gjmvq Sudden Death in Athletes
- Hypertrophic Cardiomyopathy Association www.4hcm.org
- American Heart Association www.heart.org

Collaborating Agencies:

American Academy of Pediatrics New Jersey Chapter

3836 Quakerbridge Road, Suite 108 Hamilton, NJ 08619 p) 609-842-0014

f) 609-842-0015

www.aapnj.org



American Heart Association

Union Street, Suite 301 Robbinsville, NJ, 08691 p) 609-208-0020 /ww.heart.org



New Jersey Department of Education

www.state.nj.us/education/ renton, NJ 08625-0500 (p) 609-292-5935 PO Box 500



Trenton, NJ 08625-0360 (p) 609-292-7837 P.O. Box 360

New Jersey Department of Health

www.state.nj.us/health

Lead Author: American Academy of Pediatrics, **New Jersey Chapter**

Written by: Initial draft by Sushma Raman Hebbar, MD & Stephen G. Rice, MD PhD

NJ Academy of Family Practice, Pediatric Cardiologists, Additional Reviewers: NJ Department of Education, American Heart Association/New Jersey Chapter, NJ Department of Health and Senior Services, New Jersey State School Nurses

Stephen G. Rice, MD; Jeffrey Rosenberg, MD, Lakota Kruse, MD, MPH; Susan Martz, EdM; Louis Teichholz, MD; Perry Weinstock, MD Christene DeWitt-Parker, MSN, CSN, RN; Revised 2014: Nancy Curry, EdM;

Sudden Cardiac Death ATHLETES he Basic Facts on CARDIAC SUDDE YOUNG

in Young Athletes

time) during or immediately after exercise

without trauma. Since the heart stops

heart function, usually (about 60% of the

result of an unexpected failure of proper

Sudden cardiac death is the

What is sudden cardiac death

in the young athlete?

done to prevent this kind of

tragedy?

What, if anything, can be and 19 is very rare.

> DEPARTMENT OF EDUCATION STATE OF NEW JERSEY

American Academy of Pediatrics DEDICATED TO THE HEALTH OF ALL CHILDREN*



Learn and Live

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES

What are the most common causes?

udden death in young athletes

between the ages of 10

by one of several cardiovascular abnormalities roo-LAY-shun). The problem is usually caused ventricular fibrillation (ven-TRICK-you-lar fib-Research suggests that the main cause is a and electrical diseases of the heart that go oss of proper heart rhythm, causing the blood to the brain and body. This is called unnoticed in healthy-appearing athletes. neart to quiver instead of pumping

also called HCM. HCM is a disease of the heart, muscle, which can cause serious heart rhythm The most common cause of sudden death in problems and blockages to blood flow. This (hi-per-TRO-fic CAR- dee-oh-my-OP-a-thee) genetic disease runs in families and usually an athlete is hypertrophic cardiomyopathy with abnormal thickening of the heart develops gradually over many years.

ultimately dies unless normal heart rhythm

is restored using an automated external

defibrillator (AED).

pumping adequately, the athlete quickly

collapses, loses consciousness, and

How common is sudden death in young

Sudden cardiac death in young athletes is

The chance of sudden death occurring to any individual high school athlete is

about one in 200,000 per year.

reported in the United States per year.

very rare. About 100 such deaths are

blood vessels are connected to arteries. This means that these the main blood vessel of the The second most likely cause is congenital abnormalities of the coronary (con-JEN-it-al) (i.e., present from birth)

commonly called "coronary artery disease," which may lead to a heart heart in an abnormal way. This differs from blockages that may occur when people get older

attack).

other sports; and in African-Americans than

in other races and ethnic groups.

common: in males than in females;

Sudden cardiac death is more

in football and basketball than in

12

Other diseases of the heart that can lead to What are the sudden death in young people include: for screening

- Myocarditis (my-oh-car-DIE-tis), an acute inflammation of the heart muscle (usually due to a virus).
- Dilated cardiomyopathy, an enlargement of the heart for unknown reasons.
- Long QT syndrome and other electrical abnormalities of the heart which cause abnormal fast heart rhythms that can also run in families.
- Marfan syndrome, an inherited disorder that affects heart valves, walls of major arteries, eyes and the skeleton. It is generally seen in unusually tall athletes, especially if being tall is not common in other family members.

Are there warning signs to watch for?

In more than a third of these sudden cardiac deaths, there were warning signs that were not reported or taken seriously. Warning signs are:

- Fainting, a seizure or convulsions during physical activity;
- Fainting or a seizure from emotional excitement, emotional distress or being startled;
- Dizziness or lightheadedness, especially during exertion;
- Chest pains, at rest or during exertion;
- Palpitations awareness of the heart beating unusually (skipping, irregular or extra beats) during athletics or during cool down periods after athletic participation;
- Fatigue or tiring more quickly than peers; or
- Being unable to keep up with friends due to shortness of breath (labored breathing)

What are the current recommendations for screening young athletes?

SUDDEN CARDIAC DEATH IN YOUNG ATHLETES

New Jersey requires all school athletes to be examined by their primary care physician ("medical home") or school physician at least once per year. The New Jersey Department of Education requires use of the specific Preparticipation Physical Examination Form (PPE).

This process begins with the parents and student-athletes answering questions about symptoms during exercise (such as chest pain, dizziness, fainting, palpitations or shortness of breath); and questions about family health history.

The primary healthcare provider needs to know if any family member died suddenly during physical activity or during a seizure. They also need to know if anyone in the family under the age of 50 had an unexplained sudden death such as drowning or car accidents. This information must be provided annually for each exam because it is so essential to identify those at risk for sudden cardiac death.

The required physical exam includes measurement of blood pressure and a careful listening examination of the heart, especially for murmurs and rhythm abnormalities. If there are no warning signs reported on the health history and no abnormalities discovered on exam, no further evaluation or testing is recommended.

Are there options privately available to screen for cardiac conditions?

Technology-based screening programs including a 12-lead electrocardiogram (ECG) and echocardiogram (ECHO) are noninvasive and painless options parents may consider in addition to the required

PPE. However, these procedures may be expensive and are not currently advised by the American Academy of Pediatrics and the American College of Cardiology unless the PPE reveals an indication for these tests. In addition to the expense, other limitations of technology-based tests include the possibility of "false positives" which leads to unnecessary stress for the student and parent or guardian as well as unnecessary restriction from athletic participation.

The United States Department of Health and Human Services offers risk assessment options under the Surgeon General's Family History Initiative available at http://www.hhs.gov/familyhistory/index.html

When should a student athlete see a heart specialist?

If the primary healthcare provider or school physician has concerns, a referral to a child heart specialist, a pediatric cardiologist, is recommended. This specialist will perform a more thorough evaluation, including an electrocardiogram (ECG), which is a graph of the electrical activity of the heart. An echocardiogram, which is an ultrasound test to allow for direct visualization of the heart structure, will likely also be done. The specialist may also order a treadmill exercise test and a monitor to enable a longer recording of the heart rhythm. None of the testing is invasive or uncomfortable.

Can sudden cardiac death be prevented just through proper screening?

A proper evaluation should find most, but not all, conditions that would cause sudden death in the athlete. This is because some diseases are difficult to uncover and may only develop later in life. Others can develop following a

normal screening evaluation, such as an infection of the heart muscle from a virus.

This is why screening evaluations and a review of the family health history need to be performed on a yearly basis by the athlete's primary healthcare provider. With proper screening and evaluation, most cases can be identified and prevented.

Why have an AED on site during sporting

The only effective treatment for ventricular fibrillation is immediate use of an automated external defibrillator (AED). An AED can restore the heart back into a normal rhythm. An AED is also life-saving for ventricular fibrillation caused by a blow to the chest over the heart (commotio cordis).

N.J.S.A. 18A:40-41a through c, known as "Janet's Law," requires that at any schoolsponsored athletic event or team practice in New Jersey public and nonpublic schools including any of grades K through 12, the following must be available:

- An AED in an unlocked location on school property within a reasonable proximity to the athletic field or gymnasium; and
- the athletic field or gymnasium; and

 A team coach, licensed athletic trainer, or
 other designated staff member if there is no
 coach or licensed athletic trainer present,
 certified in cardiopulmonary resuscitation
 (CPR) and the use of the AED; or
 - A State-certified emergency services provider or other certified first responder.

The American Academy of Pediatrics recommends the AED should be placed in central location that is accessible and ideally no more than a 1 to 1½ minute walk from any location and that a call is made to activate 911 emergency system while the AED is being



Sudden Cardiac Death Pamphlet Sign-Off Sheet

Name of School District:
Name of Local School:
I/We acknowledge that we received and reviewed the Sudden Cardiac Death in Young Athletes pamphlet.
Student Signature:
Parent or Guardian Signature:
Date: